

# Plasma Assisted Combustion Mechanism for Small Hydrocarbons

**Andrey Starikovskiy  
Nickolay Aleksandrov**




**PRINCETON  
University**



<b>Report Documentation Page</b>			Form Approved OMB No. 0704-0188					
<p>Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p>								
1. REPORT DATE <b>2015</b>	2. REPORT TYPE	3. DATES COVERED <b>00-00-2015 to 00-00-2015</b>						
<b>4. TITLE AND SUBTITLE</b> <b>Plasma Assisted Combustion Mechanism for Small Hydrocarbons</b>			5a. CONTRACT NUMBER					
			5b. GRANT NUMBER					
			5c. PROGRAM ELEMENT NUMBER					
<b>6. AUTHOR(S)</b>			5d. PROJECT NUMBER					
			5e. TASK NUMBER					
			5f. WORK UNIT NUMBER					
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> <b>Princeton University, Princeton, NJ, 08544</b>			8. PERFORMING ORGANIZATION REPORT NUMBER					
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>			10. SPONSOR/MONITOR'S ACRONYM(S)					
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)					
<b>12. DISTRIBUTION/AVAILABILITY STATEMENT</b> <b>Approved for public release; distribution unlimited</b>								
<b>13. SUPPLEMENTARY NOTES</b> <b>AFOSR MURI 2015 Fifth Year Review Meeting</b>								
<b>14. ABSTRACT</b>								
<b>15. SUBJECT TERMS</b>								
<b>16. SECURITY CLASSIFICATION OF:</b> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33.33%; padding: 2px;">a. REPORT <b>unclassified</b></td> <td style="width: 33.33%; padding: 2px;">b. ABSTRACT <b>unclassified</b></td> <td style="width: 33.33%; padding: 2px;">c. THIS PAGE <b>unclassified</b></td> </tr> </table>			a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	<b>17. LIMITATION OF ABSTRACT</b> <b>Same as Report (SAR)</b>	<b>18. NUMBER OF PAGES</b> <b>45</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>						

# Ignition, Combustion and Flame Control by Nonequilibrium Plasma




1814 – Brande. Flame/Field Interaction

W.T. Brande. Phil.Trans.Roy.Soc., 1814, 104, 51.



In 1860 Étienne Lenoir used an electric spark plug in his gas engine, the first internal combustion piston engine and is generally credited with the invention of the spark plug



# Propulsion Efficiency and Operating Regimes for Variety of Flight Systems





(a)





(b)



(c)




# Lean Ignition for Gas IC Engines



Physics of Nonequilibrium Systems Laboratory

- Regular spark plugs  $\lambda < 1.4$
- Regular spark plugs with thin (Iridium/Platinum) electrodes  $\lambda < 1.6$
- RF, “plasma”, etc. plugs  $\lambda < 1.8$





# GTE Lean Regimes


$T = 700 - 1300 \text{ K}$

$P = 20 - 30 \text{ atm}$

$W = 10 - 1000 \text{ MW}$



# Decreasing of Ignition Delay Time - 1994



# Kinetic Model: Previous Versions


**D.V.Zatsepin, S.M.Starikovskaia, A.Yu.Starikovskii** *Hydrogen oxidation in a stoichiometric hydrogen-air mixtures in the fast ionization wave.* Combust. Theory Modeling, 2001. V.5 pp.97-129.

**N.A.Popov.** *Effect of a Pulsed High-Current Discharge on Hydrogen–Air Mixtures.* Plasma Physics Reports, 2008, Vol. 34, No. 5, pp. 376–391.

**I.N. Kosarev, N.L. Aleksandrov, S.V. Kindysheva, S.M. Starikovskaia , A.Yu. Starikovskii.** *Kinetics of ignition of saturated hydrocarbons by nonequilibrium plasma: C<sub>2</sub>H<sub>6</sub>- to C<sub>5</sub>H<sub>12</sub>-containing mixtures.* Combustion and Flame 156 (2009) 221–233


**A.Starikovskiy, N.Aleksandrov.** *Plasma-assisted ignition and combustion.* Progress in Energy and Combustion Science 39 (2013) 61-110

# Predictive Modeling: Key to Applications




# Princeton Plasma Combustion Kinetics

## Major Pathways




# Princeton Plasma Combustion Kinetics

## *Major Pathways*




# Princeton Plasma Combustion Kinetics

## Major Pathways




# Cross-sections




# Cross-sections Available

Atmospheric	Saturated	Unsaturated	Oxygenated	Isomers
N2	CH4	C2H2	CO	iso-butane
O2	C2H6	C2H4	CH3OH	iso-propane
CO2	C3H8	C3H6	C2H5OH	neo-pentane
H2O	C4H10		CH3OCH3 DME	
O3	C5H12			
Ar	H2			
N2O				


# PAC Kinetic Mechanism




# PAC Pathways: C<sub>2</sub>H<sub>4</sub>-air




# PAC Pathways: C<sub>2</sub>H<sub>4</sub>-air




# PAC Pathways: C<sub>2</sub>H<sub>4</sub>-air




Pathway analysis in ignition process at the time of T=1360K  
 for  $\Phi = 1$ , Ar=92%, P=2.1atm and initial T=1350K. Konnov, 2014




(LLNL nButane mechanism)



(USC mechanism)




(Konnov mechanism)




(UCSD mechanism)

# Where PAC Experimental Data is Available



# Avalanche to Streamer Transition in Uniform Electric Field (air, 1 bar, 300 K, 1 cm, various E/n)




# Princeton Plasma Combustion Kinetics


## Mechanism Validation

**T = 300 K**


**Slow Oxidation of H<sub>2</sub>, C1-C10**  
P = 1-10 Torr



**Fast Gas Heating Mechanism.**  
N<sub>2</sub>-O<sub>2</sub> mixtures P = 0.2 – 1 atm




**T = 300 - 800 K**





**Oxidation Chains Development**  
in Lean H<sub>2</sub>, CO, C1-C4 - Air  
Mixtures. P = 1 atm

**T = 800 – 1700 K**


**Ignition Delay Time Reduction.**  
H<sub>2</sub>, C1-C5, C<sub>2</sub>H<sub>2</sub>, C<sub>2</sub>H<sub>5</sub>OH – O<sub>2</sub>-Ar  
Mixtures. P = 0.3-0.5 atm



**Sensitivity Analysis for Discharge and Combustion Stages**



# SDBD Discharge and Fast Heating



*Gate = 0.5 ns*

*Time shift between frames is 1 ns*

*The movie duration is 41 ns*



*Impulse Parameters*

$V = 14 \text{ kV}$


$t_{1/2} = 20 \text{ ns}$

*Frequency = 1 kHz*

*Velocity = 0.4 mm/ns*



# Potential Energy Curves of Molecular Hydrogen




$\text{H}_2(b^3\Sigma_u)$ , 8.9 eV  
 $\sigma_{\max} = 0.33 \text{ A}^2$  (17 eV)


$\text{H}_2(a^3\Sigma_g)$ , 11.8 eV  
 $\sigma_{\max} = 0.12 \text{ A}^2$  (15 eV)

$\text{H}_2(B^1\Sigma_u)$ , 11.3 eV  
 $\sigma_{\max} = 0.48 \text{ A}^2$  (40 eV)


$\text{H}_2(C^1\Pi_u)$ , 12.4 eV  
 $\sigma_{\max} = 0.40 \text{ A}^2$  (40 eV)



# Potential Energy Curves of Molecular Oxygen



# Potential Energy Curves of Molecular Nitrogen



$\text{N}_2(A^3\Sigma_u^+)$ , 6.2 eV  
 $\sigma_{\max} = 0.08 \text{ \AA}^2$  (10 eV)


$\text{N}_2(B^3\Pi_g)$ , 7.35 eV  
 $\sigma_{\max} = 0.20 \text{ \AA}^2$  (12 eV)

$\text{N}_2(C^3\Pi_u)$ , 11.03 eV  
 $\sigma_{\max} = 0.98 \text{ \AA}^2$  (14 eV)

# Major Channels of Hot Atoms Production



# Chain Initiation/Branching Reactions




$$k = 1.6 \times 10^{-10} \times \exp(-7470/T) \text{ cm}^3/\text{s}$$

$$k(300) = 2.5 \times 10^{-21} \text{ cm}^3/\text{s}$$

$$k(\text{hot}) = 1.6 \times 10^{-10} \text{ cm}^3/\text{s}$$

$$k(300, 1 \text{ atm}) = 1.6 \times 10^{-12} \text{ cm}^3/\text{s} \quad T_{\text{crit}} \sim T_{\text{autoignition}}$$



$$k = 8.5 \times 10^{-20} \times T^{2.67} \times \exp(-3160/T) \text{ cm}^3/\text{s}$$


$$k(300) = 9.3 \times 10^{-18} \text{ cm}^3/\text{s}$$

$$k(\text{hot}) = 1.5 \times 10^{-10} \text{ cm}^3/\text{s}$$


$$k(^1D) = 1.1 \times 10^{-10} \text{ cm}^3/\text{s}$$





$$k(300, 1 \text{ atm}) = 2.2 \times 10^{-14} \text{ cm}^3/\text{s} \quad T_{\text{crit}} \sim 650K$$



$$k \sim 2m/M \quad k_{gk} \sim 1.6 \times 10^{-10} \text{ cm}^3/\text{s}$$



$$k \sim 2m/M \quad k_{gk} \sim 1.3 \times 10^{-10} \text{ cm}^3/\text{s}$$




$$k = 2.6 \times 10^{-11} \text{ cm}^3/\text{s} \quad (M = O_2)$$



$$k = 1.3 \times 10^{-11} \text{ cm}^3/\text{s} \quad (M = N_2)$$

$$k = 5.2 \times 10^{-11} \text{ cm}^3/\text{s} \quad (M = H_2)$$

# Radicals Production Increase in Cold H<sub>2</sub>-Air Mixture Due to Hot Atoms Formation





# Mechanism of Fast Heating in Discharge Plasmas (high E/N)



High ( $> 200 \text{ Td}$ ) E/N:  
electron-ion and  
ion-ion  
recombination  
kinetics

# Fractional Electron Power Transferred Into Heat in $N_2:O_2$ Mixtures

$$E/N = 10^3 \text{ Td}$$



Oxygen is required  
for efficient fast heating!



...

# Experimental Setup

Current Electric Gauge

Shunt

Discharge Tube

Pressure Gauge

Pulse Generator

PS


Pump

PS  
PS




Mono-chro-mator

PS


PM




# Hexane Oxidation by Pulsed Nanosecond Discharge



# Hydrocarbon Oxidation Efficiency for $C_1-C_6$ / $O_2$ / Air Mixtures






# Calculated and Measured Times of Oxidation



# Chemical Reactions with Excited Reagents




$AB(v) + C = A + BC(w)$   
 Rate constant from  
 modified  $\alpha$ -model  
 (Starikovskii, Lashin 1996)




$$(k_{R1}/k_{R2})_{\text{exp}} = 2600 \text{ (O'Neal, Benson 1973); } (k_{R1}/k_{R2})_{\text{theor}} = 2750$$

# Kinetics. Influence of Vibrations


Distribution Of Vibrational-Excited Components





# Influence of Vibrational Excitation on Low-Temperature Kinetics



CH4-air





C4H10-air



Experiments: L Wu, J Lane, N P Cernansky, D L Miller, A A Fridman, A Yu Starikovskiy, *Proc. of Comb. Inst.*, 2010



Modelling: D Levko, A I Schedrin, V V Naumov, S Starikovskaya, 2010

# Influence of Vibrational Excitation on Low-Temperature Kinetics: $\text{H}_2\text{O}_2$ Decomposition







Measured and calculated OH decay time.  $P = 1 \text{ atm.}$   
a)  $3\%\text{H}_2 + \text{air}$ ; b)  $0.3\%\text{C}_4\text{H}_{10} + \text{air}$ .

# Plasma Shock Tube




# Discharge Dynamics




# Active Particle Production – Discharge Phase

$U_s = 943.6 \text{ m/s}$ ;  $P_0 = 17 \text{ Torr}$ ;  $P_5 = 1.04 \text{ atm}$ ;  $T_5 = 1525 \text{ K}$




# Plasma Ignition Sensitivity


**0.1 eV/mol**




## Princeton Plasma Combustion Kinetics Major Pathways




## Princeton Plasma Combustion Kinetics Major Pathways




## Princeton Plasma Combustion Kinetics Major Pathways



## Princeton Plasma Combustion Kinetics Mechanism Validation



# Other Major Results




## 3 NEW MECHANISMS:

- Radicals Production Increase Due to Translationally Hot Atoms Formation
- Mechanism of Fast Heating in Plasmas at high E/n
- Vibrational Decomposition of Peroxides (HO<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>, etc)


## EXPERIMENTAL DATABASE:

- Plasma Ignition Delay Time database for H<sub>2</sub>, C1-C5, acetylene, ethylene, ethanol



**The work was supported by**

**AFOSR  
Technical Monitor  
Dr Chiping Li**



**PRINCETON  
University**